




Introduction to Robot Intelligence [Spring 2023]

Inverse Kinematics (Part 2)

March 7, 2023

Lerrel Pinto

FK vs. IK



Solving IK for simple mechanisms

Two linked planar robot




q_1 ? q_2 ?
 $x_{des} = l_1 c_1 + l_2 c_{12}$
 $y_{des} = l_1 s_1 + l_2 s_{12}$
 $\checkmark x_{des} = \checkmark l_1 \checkmark c_1 + \checkmark l_2 \checkmark c_{12} - \checkmark l_2 \checkmark s_1 \checkmark s_2$
 $x_{des} = (l_1 + l_2 c_2) c_1 - l_2 s_2 s_1$ — (3)
 $y_{des} = l_1 s_1 + l_2 s_1 c_2 + l_2 s_2 c_1$
 $y_{des} = (l_1 + l_2 c_2) s_1 + l_2 s_2 c_1$ — (4)

$\cos(A+B) = \cos A \cos B - \sin A \sin B$
 $\sin(A+B) = \sin A \cos B + \sin B \cos A$

$q_1 = \text{atan2}(c_1, s_1)$

$A \begin{bmatrix} l_1 + l_2 c_2 & -l_2 s_2 \\ l_2 s_2 & l_1 + l_2 c_2 \end{bmatrix} \begin{bmatrix} c_1 \\ s_1 \end{bmatrix} = \begin{bmatrix} x_{des} \\ y_{des} \end{bmatrix} \Rightarrow \begin{bmatrix} c_1 \\ s_1 \end{bmatrix} = A^{-1} \begin{bmatrix} x_{des} \\ y_{des} \end{bmatrix}$

Two linked planar robot




$x_{des} = l_1 \cos q_1 + l_2 \cos(q_1 + q_2)$ — (1)
 $y_{des} = l_1 \sin q_1 + l_2 \sin(q_1 + q_2)$ — (2)

$x_{des}^2 + y_{des}^2 = l_1^2 (s_1^2 + c_1^2) + l_2^2 (c_{12}^2 + s_{12}^2) + 2 l_1 l_2 c_1 c_{12} + 2 l_1 l_2 s_1 s_{12}$
 $= l_1^2 + l_2^2 + 2 l_1 l_2 (c_1 c_{12} + s_1 s_{12})$
 $= l_1^2 + l_2^2 + 2 l_1 l_2 (\cos q_1 \cos(q_1 + q_2) + \sin q_1 \sin(q_1 + q_2))$

2 solutions for q_2


$x_{des}^2 + y_{des}^2 = l_1^2 + l_2^2 + 2 l_1 l_2 c_2$
 $\Rightarrow \cos q_2 = \frac{x^2 + y^2 - l_1^2 - l_2^2}{2 l_1 l_2}$

$\sin q_2 = \pm \sqrt{1 - \cos^2 q_2}$
 $q_2 = \text{atan2}(\cos q_2, \sin q_2)$




Solution 2b: Numerical Methods (Gradient Descent)

New Concept – The Jacobian



$$\frac{db}{da}$$



$$\frac{d\vec{b}}{da} = \begin{bmatrix} \frac{db_1}{da} \\ \frac{db_2}{da} \\ \vdots \\ \frac{db_n}{da} \end{bmatrix}$$

$$\vec{a} = [a_1, a_2, \dots, a_m] \quad b$$


$$\frac{db}{d\vec{a}} = \left[\frac{\partial b}{\partial a_1} \quad \frac{\partial b}{\partial a_2} \quad \dots \quad \frac{\partial b}{\partial a_m} \right]$$

$$\vec{a} = [a_1, a_2, \dots, a_m] \quad \vec{b} = [b_1, b_2, \dots, b_n]$$

$$\frac{d\vec{b}}{d\vec{a}} = \begin{bmatrix} \frac{\partial b_1}{\partial a_1} & \frac{\partial b_1}{\partial a_2} & \dots & \frac{\partial b_1}{\partial a_m} \\ \frac{\partial b_2}{\partial a_1} & \frac{\partial b_2}{\partial a_2} & \dots & \frac{\partial b_2}{\partial a_m} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial b_n}{\partial a_1} & \frac{\partial b_n}{\partial a_2} & \dots & \frac{\partial b_n}{\partial a_m} \end{bmatrix}_{n \times m}$$

$$= \underline{\underline{J(\vec{b}, \vec{a})}}$$

A robot's Jacobian




$$\vec{q} = [q_1, q_2, \dots, q_m]$$

$$\vec{x}(\vec{q}_0 + \Delta \vec{q}) = \vec{x}(\vec{q}) + \underline{J} \Delta \vec{q}$$

$x \rightarrow n-D$ var
 $q \rightarrow m-D$ var

$$\underline{J}(\vec{x}, \vec{q})$$


$$\frac{d\vec{x}}{d\vec{q}}$$



one D variable $a_0 + \Delta a$

$$b(a_0 + \Delta a) = b_0 + \frac{db}{da} \cdot \Delta a$$

A robot's Jacobian




$$\vec{q} = [q_1, q_2]$$
$$\vec{x} = [a, b]$$
A hand-drawn coordinate system with a vertical axis labeled a and a horizontal axis labeled b .

Image credits: Kris Hauser.

A robot's Jacobian




Image credits: Kris Hauser.

A robot's Jacobian




Image credits: Kris Hauser.

A robot's Jacobian




Figure 5. The Jacobian relates joint space velocities to Cartesian velocities. The first column of the Jacobian gives the velocity at which the end effector would move if the first joint moves at unit speed, and the second column gives its velocity if the second joint moves at unit velocity. The linear combination of Jacobian columns, weighted by joint speed, gives the overall Cartesian velocity.

Image credits: Kris Hauser.

Recall: Gradient Descent Algorithm with multiple params

Recall: Gradient Descent Algorithm with multiple params

$\vec{a}$

- Given: cost / loss/ objective function $f(\vec{\theta})$. Where $\vec{\theta} \in \mathbb{R}^d$.


- Goal: find $\vec{\theta}^*$ such that $f(\vec{\theta}^*) = \min_{\vec{\theta}} f(\vec{\theta})$.

→ • Gradient descent solution:

- • Start from initial guess $\vec{\theta}^0$ and learning rate α

- • Update $\vec{\theta}^{i+1} \leftarrow \vec{\theta}^i - \alpha \nabla f(\vec{\theta})$

- • Repeat until change in θ is small, or maximum number of steps reached.




$$\frac{d}{dx} f^2(y) = 2 f(y) \cdot \frac{dy}{dx}$$

$$f(g(x))' = f'(g(x)) g'(x)$$

Gradient Descent for IK

Vanilla Gradient Descent



$$\vec{x} \leftarrow \text{F.K.}(\vec{q}) \mid \vec{x}(\vec{q})$$

$$\min. f(\vec{q}) = \|\vec{x}(\vec{q}) - \vec{d}\|^2$$

$$\|\vec{a}\|^2 = \vec{a}^T \vec{a}$$

Objective

→ Initial $\vec{q}_0$

→ Update $\vec{q}_1 \leftarrow \vec{q}_0 - \alpha \nabla_{\vec{q}} f(\vec{q})$

→ Repeat until convergence

$$f(\vec{q}) = (\vec{x}(\vec{q}) - \vec{d})^T (\vec{x}(\vec{q}) - \vec{d})$$

$$\nabla_{\vec{q}} f(\vec{q}) = 2 \left(\frac{d\vec{x}}{d\vec{q}} \right) (\vec{x}(\vec{q}) - \vec{d})$$

$$= 2 \underline{J^T} (\vec{x}(\vec{q}) - \vec{d})$$

$$= 2 \begin{bmatrix} \frac{\partial x_1}{\partial q_1} & \frac{\partial x_2}{\partial q_1} & \dots & \frac{\partial x_n}{\partial q_1} \\ \frac{\partial x_1}{\partial q_m} & \frac{\partial x_2}{\partial q_m} & \dots & \frac{\partial x_n}{\partial q_m} \end{bmatrix} \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_n \end{bmatrix}$$

Gradient Descent for IK

$$Ax = b \quad A^{-1}$$

$$x = \underline{\underline{A^{-1}b}}$$

Jacobian-pseudoinverse gradient descent

$$J(\vec{x}, \vec{q})$$

$$\Delta x = \underset{n \times m}{J} \Delta q$$


$$\frac{db}{da}$$

$$\Delta b = \frac{db}{da} \Delta a$$

$$\Delta x_d = J \Delta q_d$$

$$\Delta q_d = J^T \Delta x_d$$

$$= (J^T J)^{-1} J^T \Delta x_d$$



(i) guess $\vec{q}_0$

(ii) compute

- $\rightarrow \vec{x}_0 = F.K(\vec{q}_0)$
- $\rightarrow (\vec{d} - \vec{x}_0)$
- $\rightarrow \propto (\vec{d} - \vec{x}_0)$

(iii) $\vec{q}_1 \leftarrow \vec{q}_0 \pm \alpha (J^T J)^{-1} J^T (\vec{d} - \vec{x}_0)$

How do I do this practically?

- Step 1: Figure out the Forward Kinematics

How do I do this practically?

- Step 1: Figure out the Forward Kinematics
- Step 2: Differentiate it!
 - Option 1: Do it by hand.
 - Option 2: Do it numerically.


How do I do this practically?

- Step 1: Figure out the Forward Kinematics
- Step 2: Differentiate it!
 - Option 1: Do it by hand.
 - Option 2: Do it numerically.
- Step 3: Choose learning rate.


How do I do this practically?

- Step 1: Figure out the Forward Kinematics
- Step 2: Differentiate it!
 - Option 1: Do it by hand.
 - Option 2: Do it numerically.
- Step 3: Choose learning rate.
- Step 4: Run optimizer.

Examples





<https://www.alanzucconi.com/>



<https://learn.foundry.com/>

Examples



<https://www.alanzucconi.com/>

Additional Reading

- <http://motion.cs.illinois.edu/RoboticSystems/Kinematics.html>
- Textbook: <http://hades.mech.northwestern.edu/images/2/2a/Park-lynch.pdf>